



Fgl2 Polyclonal Antibody

Catalog No	BYab-13927
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	FGL2
Protein Name	Fibroleukin
Immunogen	The antiserum was produced against synthesized peptide derived from human Fgl2. AA range:38-87
Specificity	Fgl2 Polyclonal Antibody detects endogenous levels of Fgl2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	FGL2; Fibroleukin; Fibrinogen-like protein 2; pT49
Observed Band	55kD
Cell Pathway	Secreted.
Tissue Specificity	Constitutively expressed in cytotoxic T-cells.
Function	function:May play a role in physiologic lymphocyte functions at mucosal sites.,similarity:Contains 1 fibrinogen C-terminal domain.,subunit:Homotetramer; disulfide-linked.,tissue specificity:Constitutively expressed in cytotoxic T-cells.,
Background	fibrinogen like 2(FGL2) Homo sapiens The protein encoded by this gene is a secreted protein that is similar to the beta- and gamma-chains of fibrinogen. The carboxyl-terminus of the encoded protein consists of the fibrinogen-related domains (FRED). The encoded protein forms a tetrameric complex which is stabilized by interchain disulfide bonds. This protein may play a role in physiologic functions at mucosal sites. [provided by RefSeq, Jul 2008],

Nanjing BYabscience technology Co.,Ltd




matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Fgl2 Polyclonal Antibody



Western blot analysis of lysate from 293 cells treated with insulin, using Fgl2 antibody.