



CD89 Polyclonal Antibody

| | |
|---------------------------|--|
| Catalog No | BYab-14046 |
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB;IHC;IF;ELISA |
| Gene Name | FCAR |
| Protein Name | Immunoglobulin alpha Fc receptor |
| Immunogen | The antiserum was produced against synthesized peptide derived from the Internal region of human FCAR. AA range:31-80 |
| Specificity | CD89 Polyclonal Antibody detects endogenous levels of CD89 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/10000.. IF 1:50-200 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | FCAR; CD89; Immunoglobulin alpha Fc receptor; IgA Fc receptor; CD89 |
| Observed Band | 32kD |
| Cell Pathway | [Isoform A.1]: Cell membrane; Single-pass type I membrane protein.; [Isoform A.2]: Cell membrane; Single-pass type I membrane protein.; [Isoform A.3]: Cell membrane; Single-pass type I membrane protein.; [Isoform B]: Secreted.; [Isoform B-delta-S2]: Secreted. |
| Tissue Specificity | Isoform A.1, isoform A.2 and isoform A.3 are differentially expressed between blood and mucosal myeloid cells. Isoform A.1, isoform A.2 and isoform A.3 are expressed in monocytes. Isoform A.1 and isoform A.2 are expressed in alveolar macrophages; however only one isoform is expressed at alveolar macrophages surfaces. |
| Function | alternative products:Additional isoforms seem to exist,function:Binds to the Fc region of immunoglobulins alpha. Mediates several functions including cytokine production.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,subunit:Associates with the Fc epsilon RI gamma 2 receptor inducing tyrosine phosphorylation of gamma 2.,tissue specificity:Differentially expressed between blood and mucosal myeloid cells. Monocytes express isoform A.1, |

Nanjing BYabscience technology Co.,Ltd



isoform A.2 and isoform A.3 while alveolar macrophages express isoform A.1 and isoform A.2; however they express only one isoform at their surfaces.,

Background

This gene is a member of the immunoglobulin gene superfamily and encodes a receptor for the Fc region of IgA. The receptor is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens. It interacts with IgA-opsonized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2008],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis of NIH-3T3 cells using CD89 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-tonsils, antibody was diluted at 1:100