



TBK1/NAK (phospho-Ser172) rabbit pAb

Catalog No	BYab-14622
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	TBK1 NAK
Protein Name	TBK1/NAK (Ser172)
Immunogen	Synthesized phospho peptide around human TBK1 and NAK (Ser172)
Specificity	This antibody detects endogenous levels of Human Mouse TBK1/NAK (phospho-Ser172)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Serine/threonine-protein kinase TBK1 (EC 2.7.11.1) (NF-kappa-B-activating kinase) (T2K) (TANK-binding kinase 1)
Observed Band	80kD
Cell Pathway	Cytoplasm . Upon mitogen stimulation or triggering of the immune system, TBK1 is recruited to the exocyst by EXOC2. .
Tissue Specificity	Ubiquitous with higher expression in testis. Expressed in the ganglion cells, nerve fiber layer and microvasculature of the retina.
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Serine/threonine protein involved in the signaling cascade converging to the activation of the transcription factor NF-kappa-B. May function as an IKK kinase, playing an essential role in the transcription of a subset of TNF-alpha-induced genes. Also mediates production of RANTES/CCL5 and interferon-beta/IFNB1. Has a pivotal role in the innate immune response. Phosphorylates Borna disease virus (BDV) P protein. Phosphorylates and activates IRF3 and IRF7 and allows their nuclear localization. This leads to production of alpha/beta interferons and the development of a cellular antiviral state. It also seems to be a central factor in the induction of the antiviral interferon

Nanjing BYabscience technology Co.,Ltd



response. Inhibition of its interaction with IRF3, due to HCV NS3 binding or BDV P protein seems to be one mechanism of inhibition of the innate immu

Background

The NF-kappa-B (NFKB) complex of proteins is inhibited by I-kappa-B (IKB) proteins, which inactivate NFKB by trapping it in the cytoplasm. Phosphorylation of serine residues on the IKB proteins by IKB kinases marks them for destruction via the ubiquitination pathway, thereby allowing activation and nuclear translocation of the NFKB complex. The protein encoded by this gene is similar to IKB kinases and can mediate NFKB activation in response to certain growth factors. [provided by RefSeq, Oct 2010],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Hafnium oxide nanoparticles coated ATR inhibitor to enhance the radiotherapy and potentiate antitumor immune response
CHEMICAL ENGINEERING JOURNAL Ruixue Liu, Chenyang Zhang, Xiaochen Wu, Chengyan Wang, Maoru Zhao, Chao Ji, Xinghua Dong, Ronghua Wang, Huanhuan Ma, Xiaochun Wang, Yan Tan, Jiangfeng Du, Zhanjun Gu
WB Mouse 4T1 cell



Hafnium oxide nanoparticles coated ATR inhibitor to enhance the radiotherapy and potentiate antitumor immune response
CHEMICAL ENGINEERING JOURNAL Ruixue Liu, Chenyang Zhang, Xiaochen Wu, Chengyan Wang, Maoru Zhao, Chao Ji, Xinghua Dong, Ronghua Wang, Huanhuan Ma, Xiaochun Wang, Yan Tan, Jiangfeng Du, Zhanjun Gu
WB Mouse 4T1 cell