





HoxA5 Polyclonal Antibody

Catalog No BYab-15761 Isotype IgG Reactivity Human; Mouse; Rat Applications WB; ELISA Gene Name HOXA5 Protein Name Homeobox protein Hox-A5 Immunogen Synthesized peptide derived from the N-terminal region of human HoxA5. Specificity HoxA5 Polyclonal Antibody detects endogenous levels of HoxA5 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary, Function developmental stage:Expressed during embryogenesis and in adult kidney, function:Sequence-specific transcription factor which is part of a devel		
Reactivity Human;Mouse;Rat Applications WB;ELISA Gene Name HOXA5 Protein Name Homeobox protein Hox-A5 Immunogen Synthesized peptide derived from the N-terminal region of human HoxA5. Specificity HoxA5 Polyclonal Antibody detects endogenous levels of HoxA5 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary. Function developmental stage:Expressed during embryogenesis and in adult kidney, function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promote	Catalog No	BYab-15761
Applications WB;ELISA Gene Name HOXA5 Protein Name Homeobox protein Hox-A5 Immunogen Synthesized peptide derived from the N-terminal region of human HoxA5. Specificity HoxA5 Polyclonal Antibody detects endogenous levels of HoxA5 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary, developmental stage:Expressed during embryogenesis and in adult kidney, function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5"-CYYNATTAT[TG]V-3; similarity-Belongs to the Antphomeobox genes are found in clusters named A, B, C, and On four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosomes.	Isotype	IgG
Protein Name Homeobox protein Hox-A5 Immunogen Synthesized peptide derived from the N-terminal region of human HoxA5. Specificity HoxA5 Polyclonal Antibody detects endogenous levels of HoxA5 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary, Function developmental stage:Expressed during embryogenesis and in adult kidney., function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with sepcific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5'-CYYNATTATTGTy-3', similarity:Belongs to the Antp homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosomes.	Reactivity	Human;Mouse;Rat
Protein Name Homeobox protein Hox-A5 Immunogen Synthesized peptide derived from the N-terminal region of human HoxA5. Specificity HoxA5 Polyclonal Antibody detects endogenous levels of HoxA5 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary, Function developmental stage:Expressed during embryogenesis and in adult kidney, function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5-CryNATTA[TG]-3-3, similarity:Belongs to the Antphomeobox family, similarity:Contains 1 homeobox DNA-binding domain., Background In ve	Applications	WB;ELISA
Immunogen Synthesized peptide derived from the N-terminal region of human HoxA5. Specificity HoxA5 Polyclonal Antibody detects endogenous levels of HoxA5 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary, Function developmental stage:Expressed during embryogenesis and in adult kidney, function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5"-C"YNATTA[TG]Y-3", similarity:Belongs to the Antphomeobox family, similarity:Contains 1 homeobox DNA-binding domain., Background In vertebrates, the genes encoding the class of transcription factors called homeobox genes ar	Gene Name	HOXA5
Specificity HoxA5 Polyclonal Antibody detects endogenous levels of HoxA5 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary, Function developmental stage:Expressed during embryogenesis and in adult kidney, function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to sown promoter. Binds specifically to the motif 5'-CYYNATTA[TG]Y-3', similarity:Belongs to the Antphomeobox family, similarity:Contains 1 homeobox DNA-binding domain. Background In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated	Protein Name	Homeobox protein Hox-A5
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary, developmental stage:Expressed during embryogenesis and in adult kidney, function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identitities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5'-CYYNATTA[TG]Y-3', similarity:Belongs to the Antphomeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosomes.	Immunogen	Synthesized peptide derived from the N-terminal region of human HoxA5.
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary, Function developmental stage:Expressed during embryogenesis and in adult kidney, function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5°-CYYNATTA[TG]Y-3°, similarity:Belongs to the Antphomeobox family, similarity:Contains 1 homeobox DNA-binding domain. Background In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosome.	Specificity	HoxA5 Polyclonal Antibody detects endogenous levels of HoxA5 protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary, developmental stage:Expressed during embryogenesis and in adult kidney., function: Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5'-CYYNATTA[TG]Y-3', similarity:Belongs to the Antp homeobox family., similarity:Contains 1 homeobox DNA-binding domain. In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosomes.	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary, Function developmental stage:Expressed during embryogenesis and in adult kidney.,function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5'-CYYNATTA[TG]Y-3'.,similarity:Belongs to the Antphomeobox family.,similarity:Contains 1 homeobox DNA-binding domain., In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosomes.	Source	Polyclonal, Rabbit,IgG
applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary, Function developmental stage:Expressed during embryogenesis and in adult kidneyfunction:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5¹-CYYNATTA[TG]Y-3¹', similarity:Belongs to the Antp homeobox family.,similarity:Contains 1 homeobox DNA-binding domain., Background In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosomes.	Purification	·
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary, Function developmental stage:Expressed during embryogenesis and in adult kidney.,function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5°-CYYNATTA[TG]Y-3",similarity:Belongs to the Antphomeobox family.,similarity:Contains 1 homeobox DNA-binding domain., Background In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosomes.	Dilution	
Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary, Function developmental stage: Expressed during embryogenesis and in adult kidney., function: Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5'-CYYNATTA[TG]Y-3'., similarity: Belongs to the Antp homeobox family., similarity: Contains 1 homeobox DNA-binding domain., Background In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosome.	Concentration	• •
Synonyms HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C Observed Band 40kD Cell Pathway Nucleus. Tissue Specificity Ovary, Function developmental stage: Expressed during embryogenesis and in adult kidney., function: Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5'-CYYNATTA[TG]Y-3'., similarity: Belongs to the Antp homeobox family., similarity: Contains 1 homeobox DNA-binding domain., In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosomes.	Purity	≥90%
Cell Pathway Nucleus. Tissue Specificity Ovary, developmental stage: Expressed during embryogenesis and in adult kidney., function: Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5'-CYYNATTA[TG]Y-3'., similarity: Belongs to the Antp homeobox family., similarity: Contains 1 homeobox DNA-binding domain., In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosomes.	Storage Stability	-20°C/1 year
Cell Pathway Nucleus. Tissue Specificity Ovary, developmental stage: Expressed during embryogenesis and in adult kidney., function: Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5'-CYYNATTA[TG]Y-3'., similarity: Belongs to the Antp homeobox family., similarity: Contains 1 homeobox DNA-binding domain., In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosomes.	Synonyms	HOXA5; HOX1C; Homeobox protein Hox-A5; Homeobox protein Hox-1C
Tissue Specificity Ovary, developmental stage: Expressed during embryogenesis and in adult kidney., function: Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5'-CYYNATTA[TG]Y-3'., similarity: Belongs to the Antp homeobox family., similarity: Contains 1 homeobox DNA-binding domain., Background In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosomes.	Observed Band	40kD
developmental stage:Expressed during embryogenesis and in adult kidney.,function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5'-CYYNATTA[TG]Y-3'.,similarity:Belongs to the Antp homeobox family.,similarity:Contains 1 homeobox DNA-binding domain., Background In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosomes.	Cell Pathway	Nucleus.
kidney.,function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5'-CYYNATTA[TG]Y-3'.,similarity:Belongs to the Antp homeobox family.,similarity:Contains 1 homeobox DNA-binding domain., Background In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosome	Tissue Specificity	Ovary,
homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosome	Function	kidney.,function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Also binds to its own promoter. Binds specifically to the motif 5'-CYYNATTA[TG]Y-3'.,similarity:Belongs to the Antp
	Background	homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosome

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	expression, morphogenesis, and differentiation. Methylation of this gene may result in the loss of its expression and, since the encoded protein upregulates the tumor suppressor p53, this protein may play an important role in tumorigenesis. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of 293T cells using HoxA5 Polyclonal Antibody diluted at 1:500 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

Nanjing BYabscience technology Co.,Ltd