



Cleaved-Ephrin-A4 (S170) rabbit pAb

| | |
|----------------------------------|---|
| Catalog No | BYab-15821 |
| Isotype | IgG |
| Reactivity | Human;Mouse |
| Applications | WB; ELISA |
| Gene Name | EFNA4 EPLG4 LERK4 |
| Protein Name | Cleaved-Ephrin-A4 (S170) |
| Immunogen | Synthesized peptide derived from human Cleaved-Ephrin-A4 (S170) |
| Specificity | This antibody detects endogenous levels of Human,Mouse Cleaved-Ephrin-A4 (S170, protein was cleaved amino acid sequence between 170-171) |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Dilution | WB 1:1000-2000 ELISA 1:5000-20000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | Ephrin-A4 (EPH-related receptor tyrosine kinase ligand 4;LERK-4) |
| Observed Band | 17kD |
| Cell Pathway | [Isoform 1]: Cell membrane; Lipid-anchor, GPI-anchor.; [Isoform 2]: Secreted . |
| Tissue Specificity | Expressed in the adult spleen, lymph node, prostate, ovary, small intestine, and colon, and in fetal heart, lung, liver and kidney. Also detected in hematopoietic cell lines. |
| Function | cell-cell signaling, |
| Background | function:May play a role in the interaction between activated B lymphocytes and dendritic cells in tonsils.,similarity:Belongs to the ephrin family.,tissue specificity:Expressed in the adult spleen, lymph node, prostate, ovary, small intestine, and colon, and in fetal heart, lung, liver and kidney. Also detected in hematopoietic cell lines., |
| matters needing attention | Avoid repeated freezing and thawing! |

Nanjing BYabscience technology Co.,Ltd



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images