



## **BMP-8** Polyclonal Antibody

| Catalog No         | BYab-15881  |
|--------------------|---|
| Isotype            | lgG   |
| Reactivity         | Human;Rat;Mouse;  |
| Applications       | IHC;IF;ELISA  |
| Gene Name          | BMP8A/BMP8B   |
| Protein Name       | Bone morphogenetic protein 8A/B   |
| Immunogen          | The antiserum was produced against synthesized peptide derived from human BMP-8. AA range:41-90   |
| Specificity        | BMP-8 Polyclonal Antibody detects endogenous levels of BMP-8 protein.   |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source             | Polyclonal, Rabbit,IgG  |
| Purification       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Dilution           | IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200  |
| Concentration      | 1 mg/ml   |
| Purity             | ≥90%  |
| Storage Stability  | -20°C/1 year  |
| Synonyms           | BMP8B; BMP8; Bone morphogenetic protein 8B; BMP-8; BMP-8B; Osteogenic protein 2; OP-2; BMP8A; Bone morphogenetic protein 8A; BMP-8A   |
| Observed Band      |   |
| Cell Pathway       | Secreted .  |
| Tissue Specificity | Hippocampus,Testis,   |
| Function           | function:Induces cartilage and bone formation. May be the osteoinductive factor<br>responsible for the phenomenon of epithelial osteogenesis. Plays a role in<br>calcium regulation and bone homeostasis.,online information:Bone<br>morphogenetic protein 8b entry,similarity:Belongs to the TGF-beta<br>family.,subunit:Homodimer; disulfide-linked.,                                 |
| Background         | This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked |
|                    | Nanjing BYabscience technology Co.,Ltd  |

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商

精准的 WB 检测服务 24H 在线服务, 欢迎咨询

|                           | homodimer. The encoded protein stimulates thermogenesis in brown adipose<br>tissue. Expression of this gene may be downregulated in pancreatic cancer. This<br>gene may have arose from a gene duplication event and its gene duplicate is also<br>present on chromosome 1. [provided by RefSeq, Jul 2016], |
|---------------------------|---|
| matters needing attention | Avoid repeated freezing and thawing!  |
| Usage suggestions         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.   |

## **Products Images**



Immunohistochemistry analysis of BMP-8 antibody in paraffin-embedded human testis tissue.

Nanjing BYabscience technology Co.,Ltd