



CRF Polyclonal Antibody

| | |
|---------------------------|--|
| Catalog No | BYab-15893 |
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | IHC;IF;ELISA |
| Gene Name | CRH |
| Protein Name | Corticoliberin |
| Immunogen | The antiserum was produced against synthesized peptide derived from human CRF. AA range:104-153 |
| Specificity | CRF Polyclonal Antibody detects endogenous levels of CRF protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | IHC: 1/100 - 1/300. ELISA: 1/10000.. IF 1:50-200 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | CRH; Corticoliberin; Corticotropin-releasing factor; CRF; Corticotropin-releasing hormone |
| Observed Band | |
| Cell Pathway | Secreted . |
| Tissue Specificity | Produced by the hypothalamus and placenta. |
| Function | function:This hormone from hypothalamus regulates the release of corticotropin from pituitary gland.,online information:Corticotropin-releasing hormone entry,similarity:Belongs to the sauvagine/corticotropin-releasing factor/urotensin I family., |
| Background | This gene encodes a member of the corticotropin-releasing factor family. The encoded preproprotein is proteolytically processed to generate the mature neuropeptide hormone. In response to stress, this hormone is secreted by the paraventricular nucleus (PVN) of the hypothalamus, binds to corticotropin releasing hormone receptors and stimulates the release of adrenocorticotrophic hormone from the pituitary gland. Marked reduction in this protein has been |

Nanjing BYabscience technology Co.,Ltd



observed in association with Alzheimer's disease. Autosomal recessive hypothalamic corticotropin deficiency has multiple and potentially fatal metabolic consequences including hypoglycemia and hepatitis. In addition to production in the hypothalamus, this protein is also synthesized in peripheral tissues, such as T lymphocytes, and is highly expressed in the placenta. In the placenta it is a marker that determines the length of gestation a


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of CRF antibody in paraffin-embedded human lung carcinoma tissue.