



Ghrelin Polyclonal Antibody

Catalog No	BYab-15918
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	GHRL
Protein Name	Appetite-regulating hormone
Immunogen	The antiserum was produced against synthesized peptide derived from human Ghrelin. AA range:47-96
Specificity	Ghrelin Polyclonal Antibody detects endogenous levels of Ghrelin protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. IF 1:100-300 Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	GHRL; MTLRP; Appetite-regulating hormone; Growth hormone secretagogue; Growth hormone-releasing peptide; Motilin-related peptide; Protein M46
Observed Band	13kD
Cell Pathway	Secreted.
Tissue Specificity	Highest level in stomach. All forms are found in serum as well. Other tissues compensate for the loss of ghrelin synthesis in the stomach following gastrectomy.
Function	function:Ghrelin is the ligand for growth hormone secretagogue receptor type 1 (GHSR). Induces the release of growth hormone from the pituitary. Has an appetite-stimulating effect, induces adiposity and stimulates gastric acid secretion. Involved in growth regulation.,function:Obestatin may be the ligand for GPR39. May have an appetite-reducing effect resulting in decreased food intake. May reduce gastric emptying activity and jejunal motility.,mass spectrometry:Ghrelin-27-C10, O-decanoylated form PubMed:12414809,mass spectrometry:Ghrelin-27-C8, O-octanoylated form PubMed:12414809,mass spectrometry:Ghrelin-28-C10, O-decanoylated form PubMed:12414809,mass

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询




	spectrometry:Ghrelin-28-C10:1, O-decenoylated form PubMed:12414809,mass spectrometry:Ghrelin-28-C8, O-octanoylated form PubMed:12414809,online information:Ghrelin entry,online information:Gut feelings - Issue 66 of January 2006,PTM:Amida
Background	This gene encodes the ghrelin-obestatin preproprotein that is cleaved to yield two peptides, ghrelin and obestatin. Ghrelin is a powerful appetite stimulant and plays an important role in energy homeostasis. Its secretion is initiated when the stomach is empty, whereupon it binds to the growth hormone secretagogue receptor in the hypothalamus which results in the secretion of growth hormone (somatotropin). Ghrelin is thought to regulate multiple activities, including hunger, reward perception via the mesolimbic pathway, gastric acid secretion, gastrointestinal motility, and pancreatic glucose-stimulated insulin secretion. It was initially proposed that obestatin plays an opposing role to ghrelin by promoting satiety and thus decreasing food intake, but this action is still debated. Recent reports suggest multiple metabolic roles for obestatin, including regulating adipocyte function
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.







Products Images




Immunofluorescence analysis of A549. 1,primary Antibody was diluted at 1:200(4°C overnight). 2, Goat Anti Rabbit IgG (H&L) - Alexa Fluor 488 Secondary antibody was diluted at 1:1000(room temperature, 50min).3, Picture B: DAPI(blue) 10min.



Western Blot analysis of various cells using Ghrelin Polyclonal Antibody




Western Blot analysis of 293 cells using Ghrelin Polyclonal Antibody




Immunohistochemical analysis of paraffin-embedded Human lung cancer. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

Nanjing BYabscience technology Co.,Ltd









Immunohistochemistry analysis of Ghrelin antibody in paraffin-embedded human lung carcinoma tissue.

Nanjing BYabscience technology Co.,Ltd





-85

-49


-34


-25

-19

Ghrelin-

Nanjing BYabscience technology Co.,Ltd







Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 监督电话: 15950492658 官方热线: 025-5229-8998