





FGF-5 Polyclonal Antibody

| Catalog No | BYab-16003 |
|--------------------|--|
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB;IHC;IF;ELISA |
| Gene Name | FGF5 |
| Protein Name | Fibroblast growth factor 5 |
| lmmunogen | The antiserum was produced against synthesized peptide derived from the C-terminal region of human FGF5. AA range:211-260 |
| Specificity | FGF-5 Polyclonal Antibody detects endogenous levels of FGF-5 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000 IF 1:50-200 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | FGF5; Fibroblast growth factor 5; FGF-5; Heparin-binding growth factor 5; HBGF-5; Smag-82 |
| Observed Band | 30kD |
| Cell Pathway | Secreted . |
| Tissue Specificity | Expressed in neonatal brain. |
| Function | developmental stage:Can transform NIH 3T3 cells.,function:Functions as an inhibitor of hair elongation by promoting progression from anagen, the growth phase of the hair follicle, into catagen the apoptosis-induced regression phase.,similarity:Belongs to the heparin-binding growth factors family.,tissue specificity:Expressed in neonatal brain., |
| Background | The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This gene was identified as an oncogene, which confers |
| | Naniina Pyahasianaa taahnalaay Ca Itd |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询




| | transforming potential when transfected into mammalian cells. Targeted disruption of the homolog of this gene in mouse resulted in the phenotype of abnormally long hair, which suggested a function as an inhibitor of hair elongation. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008], |
|---------------------------|--|
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |







Products Images



Western Blot analysis of mouse liver cells using FGF-5 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100



Western blot analysis of lysate from mouse liver cells, using FGF5 Antibody.

Nanjing BYabscience technology Co.,Ltd