



## **VEGF-C Polyclonal Antibody**

BYab-16012
DTab-10012
lgG
Human;Mouse;Rat
WB;IHC;IF;ELISA
VEGFC
Vascular endothelial growth factor C
The antiserum was produced against synthesized peptide derived from the Internal region of human VEGFC. AA range:91-140
VEGF-C Polyclonal Antibody detects endogenous levels of VEGF-C protein.
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
WB: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000 IF 1:50-200
1 mg/ml
≥90%
-20°C/1 year
VEGFC; Vascular endothelial growth factor C; VEGF-C; Flt4 ligand; Flt4-L; Vascular endothelial growth factor-related protein; VRP
42kD
42kD Secreted.

## Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商

精准的 WB 检测服务 24H 在线服务,欢迎咨询

Before secretion, a cleavage occurs between Arg-227 and Ser-228 producing an
heterotetramer. The next extracellular step of the processing removes the N-terminal propeptide. Finally the mature VEGF-C is compos
The protein encoded by this gene is a member of the platelet-derived growth factor/vascular endothelial growth factor (PDGF/VEGF) family. The encoded protein promotes angiogenesis and endothelial cell growth, and can also affect the permeability of blood vessels. The proprotein is further cleaved into a fully processed form that can bind and activate VEGFR-2 and VEGFR-3 receptors. [provided by RefSeq, Apr 2014],
Avoid repeated freezing and thawing!
This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询




## **Products Images**



Zhou, X. U., et al. "miR-128 downregulation promotes growth and metastasis of bladder cancer cells and involves VEGF-C upregulation." Oncology letters 10.5 (2015): 3183-3190.



Western Blot analysis of L929 cells using VEGF-C Polyclonal Antibody. Antibody was diluted at 1:2000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded mouse-heart, antibody was diluted at 1:100

Nanjing BYabscience technology Co.,Ltd

官方热线: 025-5229-8998




国内优质抗体供应商 精准的 WB 检测服务

24H 在线服务, 欢迎咨询



Immunohistochemical analysis of paraffin-embedded mouse-heart, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human Liver. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).




网址: www.njbybio.com

官方热线: 025-5229-8998









Nanjing BYabscience technology Co.,Ltd

网址:www.njbybio.com 官方热线:025-5229-8998 监督电话:15950492658









Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Nanjing BYabscience technology Co.,Ltd

网址:www.njbybio.com 官方热线:025-5229-8998 监督电话:15950492658