



CLIC3 Polyclonal Antibody

| | |
|---------------------------|--|
| Catalog No | BYab-16404 |
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB;ELISA |
| Gene Name | CLIC3 |
| Protein Name | Chloride intracellular channel protein 3 |
| Immunogen | The antiserum was produced against synthesized peptide derived from human CLIC3. AA range:21-70 |
| Specificity | CLIC3 Polyclonal Antibody detects endogenous levels of CLIC3 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | CLIC3; Chloride intracellular channel protein 3 |
| Observed Band | 27kD |
| Cell Pathway | Nucleus. Membrane; Single-pass membrane protein. Cytoplasm. Predominantly nuclear. Some protein was found in the cytoplasm. Exists both as soluble cytoplasmic protein and as membrane protein with probably a single transmembrane domain (By similarity). . |
| Tissue Specificity | Detected in placenta (at protein level). Widely expressed. High expression is found in placenta followed by lung and heart. Low expression in skeletal muscle, kidney and pancreas. |
| Function | domain:Members of this family may change from a globular, soluble state to a state where the N-terminal domain is inserted into the membrane and functions as chloride channel. A conformation change of the N-terminal domain is thought to expose hydrophobic surfaces that trigger membrane insertion.,function:Can insert into membranes and form chloride ion channels. May participate in cellular growth control.,similarity:Belongs to the chloride channel CLIC family.,similarity:Contains 1 GST C-terminal domain.,similarity:Contains 1 GST N-terminal |

Nanjing BYabscience technology Co.,Ltd



domain.,subcellular location:Predominantly nuclear. Some protein was found in the cytoplasm. Exists both as soluble cytoplasmic protein and as membrane protein with probably a single transmembrane domain.,subunit:Associated with the C-terminal of ERK7.,tissue specificity:Detected in placenta (at protein level). Widely expressed. High expression is fou

Background

chloride intracellular channel 3(CLIC3) Homo sapiens Chloride channels are a diverse group of proteins that regulate fundamental cellular processes including stabilization of cell membrane potential, transepithelial transport, maintenance of intracellular pH, and regulation of cell volume. Chloride intracellular channel 3 is a member of the p64 family and is predominantly localized in the nucleus and stimulates chloride ion channel activity. In addition, this protein may participate in cellular growth control, based on its association with ERK7, a member of the MAP kinase family. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



K562

Western blot analysis of lysates from K562 cells, using CLIC3 Antibody. The lane on the right is blocked with the synthesized peptide.



CLIC3



Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658