



Na⁺ CP type IV α Polyclonal Antibody

Catalog No	BYab-16472
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC
Gene Name	SCN4A
Protein Name	Sodium channel protein type 4 subunit alpha
Immunogen	The antiserum was produced against synthesized peptide derived from human SCN4A. AA range:431-480
Specificity	Na ⁺ CP type IV α Polyclonal Antibody detects endogenous levels of Na ⁺ CP type IV α protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	SCN4A; Sodium channel protein type 4 subunit alpha; SkM1; Sodium channel protein skeletal muscle subunit alpha; Sodium channel protein type IV subunit alpha; Voltage-gated sodium channel subunit alpha Nav1.4
Observed Band	200kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein .
Tissue Specificity	Skeletal muscle,
Function	disease:Defects in SCN4A are a cause of periodic paralysis hypokalemic (HOKPP) [MIM:170400]; also designated HYPOPP. HOKPP is an autosomal dominant disorder manifested by episodic flaccid generalized muscle weakness associated with falls of serum potassium levels.,disease:Defects in SCN4A are the cause of a congenital myasthenic syndrome due to mutation in SCNA4 (CMSSCNA4) [MIM:603967]. CMSSCNA4 is a congenital myasthenic syndrome associated with fatigable generalized weakness and recurrent attacks of respiratory and bulbar paralysis since birth. The fatigable weakness involves lid-elevator, external ocular, facial, limb and truncal muscles and an decremental

Nanjing BYabscience technology Co.,Ltd



response of the compound muscle action potential on repetitive stimulation.,disease:Defects in SCN4A are the cause of myotonia SCN4A-related (MYOSCN4A) [MIM:608390]. Myotonia is characterized by sustained muscle tensing that prevent

Background

Voltage-gated sodium channels are transmembrane glycoprotein complexes composed of a large alpha subunit with 24 transmembrane domains and one or more regulatory beta subunits. They are responsible for the generation and propagation of action potentials in neurons and muscle. This gene encodes one member of the sodium channel alpha subunit gene family. It is expressed in skeletal muscle, and mutations in this gene have been linked to several myotonia and periodic paralysis disorders. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images




K5


Western blot analysis of SCN4A Antibody. The lane on the right is blocked with the SCN4A peptide.

SCN4A





Western blot analysis of the lysates from COLO205 cells using SCN4A antibody.



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).