



CTR2 rabbit pAb

Catalog No	BYab-17135
Isotype	IgG
Reactivity	Human, Mouse,Rat
Applications	IHC,WB
Gene Name	SLC7A2 ATRC2 CAT2
Protein Name	Low affinity cationic amino acid transporter 2 (CAT-2) (CAT2) (Solute carrier family 7 member 2)
Immunogen	Synthesized peptide derived from human C-terminal CTR2
Specificity	This antibody detects endogenous levels of CTR2 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Rabbit,polyclonal
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000 IHC 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Low affinity cationic amino acid transporter 2 (CAT-2) (CAT2) (Solute carrier family 7 member 2)
Observed Band	
Cell Pathway	Cell membrane ; Multi-pass membrane protein .
Tissue Specificity	Expressed at high levels in the skeletal muscle, placenta and ovary. Expressed at intermediate levels in the liver and pancreas and at low levels in the kidney and heart.
Function	Functions as permease involved in the transport of the cationic amino acids (arginine, lysine and ornithine); the affinity for its substrates differs between isoforms created by alternative splicing. Isoform 1 functions as permease that mediates the transport of the cationic amino acids (arginine, lysine and ornithine), and it has much higher affinity for arginine than isoform 2. Isoform 2 functions as low-affinity, high capacity permease involved in the transport of the cationic amino acids (arginine, lysine and ornithine) . May play a role in classical or alternative activation of macrophages via its role in arginine transport.

Nanjing BYabscience technology Co.,Ltd



Background

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images