



B-Myb (Phospho Thr494/497) Rabbit pAb

Catalog No	BYab-17158
Isotype	IgG
Reactivity	Human, Mouse
Applications	IHC, WB
Gene Name	MYBL2 BMYB
Protein Name	Myb-related protein B (B-Myb) (Myb-like protein 2)
Immunogen	Synthesized peptide derived from human B-Myb (Phospho Thr494/497)
Specificity	This antibody detects endogenous levels of B-Myb (Phospho Thr494/497) Rabbit pAb at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Rabbit, polyclonal
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000 IHC 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Myb-related protein B (B-Myb) (Myb-like protein 2)
Observed Band	80kD
Cell Pathway	Nucleus.
Tissue Specificity	
Function	function: Transcription factor involved in the regulation of cell survival, proliferation, and differentiation. Transactivates the expression of the CLU gene. PTM: Phosphorylated by cyclin A/CDK2 during S-phase. Phosphorylation at Thr-520 is probably involved in transcriptional activity. similarity: Contains 3 HTH myb-type DNA-binding domains. subunit: Component of the DREAM complex (also named LINC complex) at least composed of E2F4, E2F5, LIN9, LIN37, LIN52, LIN54, MYBL1, MYBL2, RBL1, RBL2, RBBP4, TFDP1 and TFDP2. The complex exists in quiescent cells where it represses cell cycle-dependent genes. It dissociates in S phase when LIN9, LIN37, LIN52 and LIN54 form a subcomplex that binds to MYBL22.

Nanjing BYabscience technology Co., Ltd



Background

MYB proto-oncogene like 2(MYBL2) Homo sapiens The protein encoded by this gene, a member of the MYB family of transcription factor genes, is a nuclear protein involved in cell cycle progression. The encoded protein is phosphorylated by cyclin A/cyclin-dependent kinase 2 during the S-phase of the cell cycle and possesses both activator and repressor activities. It has been shown to activate the cell division cycle 2, cyclin D1, and insulin-like growth factor-binding protein 5 genes. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of 1 HeLa cell, 2 LPS 100ng/mL 30min treated ,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000