



## P2RX7 rabbit pAb

Catalog No	BYab-17274
Isotype	IgG
Reactivity	Human, Mouse,Rat
Applications	IHC,WB
Gene Name	P2RX7
Protein Name	P2X purinoceptor 7 (P2X7) (ATP receptor) (P2Z receptor) (Purinergic receptor)
Immunogen	Synthesized peptide derived from human C-ternal P2RX7
Specificity	This antibody detects endogenous levels of P2RX7 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Rabbit,polyclonal
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000 IHC 1:50-200
Concentration	1 mg/ml
	>000/
Purity	≥90%
Purity Storage Stability	-20°C/1 year
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
Storage Stability Synonyms Observed Band	-20°C/1 year P2X purinoceptor 7 (P2X7) (ATP receptor) (P2Z receptor) (Purinergic receptor)
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year P2X purinoceptor 7 (P2X7) (ATP receptor) (P2Z receptor) (Purinergic receptor)  Cell membrane ; Multi-pass membrane protein .
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year  P2X purinoceptor 7 (P2X7) (ATP receptor) (P2Z receptor) (Purinergic receptor)  Cell membrane; Multi-pass membrane protein.  Widely expressed with highest levels in brain and immune tissues.  Receptor for ATP that acts as a ligand-gated ion channel. Responsible for ATP-dependent lysis of macrophages through the formation of membrane pores permeable to large molecules. Could function in both fast synaptic transmission and the ATP-mediated lysis of antigen-presenting cells. In the absence of its natural ligand, ATP, functions as a scavenger receptor in the recognition and
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year  P2X purinoceptor 7 (P2X7) (ATP receptor) (P2Z receptor) (Purinergic receptor)  Cell membrane; Multi-pass membrane protein.  Widely expressed with highest levels in brain and immune tissues.  Receptor for ATP that acts as a ligand-gated ion channel. Responsible for ATP-dependent lysis of macrophages through the formation of membrane pores permeable to large molecules. Could function in both fast synaptic transmission and the ATP-mediated lysis of antigen-presenting cells. In the absence of its natural ligand, ATP, functions as a scavenger receptor in the recognition and

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



ı	Isaq	<b>P</b> S	ua	ae	stic	าทร
u	Jau	C 3	uч	uc.	วแง	ノロコ

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images						

Nanjing BYabscience technology Co.,Ltd