



c-Myc Tag Rabbit pAb

Catalog No	BYab- 17766
Isotype	IgG
Reactivity	Species-independent
Applications	WB
Gene Name	-
Alternative Names	c-Myc; Myc tag; Myc protein; Myc epitope tag; Myc2
Research Field	Tags & Cell Markers
Product Categories	Tag antibody
Host	Rabbit
Molecular Weight	Refer to figures
Clonality	Polyclonal Antibody
Clonality No.	-
Dilution	WB: 1/1000-1/10000
Immunogen	Synthetic peptide EQKLISEEDL coupled to KLH.
Purification	Affinity Chromatography
Conjugation	Unconjugated
Modification	Unmodified
Form	Liquid
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Concentration	1 mg/ml
Purity	≥90%
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Nanjing BYabscience technology Co.,Ltd



Background

Myc tag is a polypeptide protein tag derived from the c-myc gene product that can be added to a protein using recombinant DNA technology. It can be used for affinity chromatography, and then used to separate recombinant, overexpressed protein from wild type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits.

Thank you for your recent purchase

Focus on your research Service for life science

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658



be used as loading control for these tissues.


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images





Thank you for your recent purchase Focus on your research Service for life science

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658