



CD163 Mouse mAb

Catalog No BYab-17785 Isotype Mouse IgG1 Reactivity Human,Mouse Applications WB,IHC-P,ELISA Gene Name CD163 Alternative Names M130; MM130; SCARI1 Research Field Immunology Product Categories Primary antibody Host Mouse Molecular Weight Calculated MW: 125 kDa Clonality Monoclonal Antibody Clonality No. 9H9-H6-W6 Dilution WB: 1/500-1/1000 IHC: 1/100-1/200 ELISA: 1/10000 Immunogen Purified recombinant fragment of human CD163 expressed in E. Coli. Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) su		
Reactivity Human,Mouse Applications WB,IHC-P,ELISA Gene Name CD163 Alternative Names M130; MM130; SCARI1 Research Field Immunology Product Categories Primary antibody Host Mouse Molecular Weight Calculated MW: 125 kDa Clonality Monoclonal Antibody Clonality Monoclonal Antibody Clonality No. 9H9-H6-W6 Dilution WB: 1/500-1/1000 IHC: 1/100-1/200 ELISA: 1/10000 Immunogen Purified recombinant fragment of human CD163 expressed in E. Coli. Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/handolobin complexs by	Catalog No	BYab-17785
Applications WB,IHC-P,ELISA Gene Name CD163 Alternative Names M130; MM130; SCARI1 Research Field Immunology Product Categories Primary antibody Host Mouse Molecular Weight Calculated MW: 125 kDa Clonality Monoclonal Antibody Clonality Monoclonal Antibody Clonality No. 9H9-H6-W6 Dilution WB: 1/500-1/1000 IHC: 1/100-1/200 ELISA: 1/10000 Immunogen Purified recombinant fragment of human CD163 expressed in E. Coli. Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involves by clearance and endocytosis of hemoglobin/handolobin complexes by	Isotype	Mouse IgG1
Alternative Names M130; MM130; SCARI1 Research Field Immunology Product Categories Primary antibody Host Mouse Molecular Weight Calculated MW: 125 kDa Clonality Monoclonal Antibody Clonality No. 9H9-H6-W6 Dilution WB: 1/500-1/1000 IHC: 1/100-1/200 ELISA: 1/10000 Immunogen Purified recombinant fragment of human CD163 expressed in E. Coli. Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemolopion/haboratoollobic norvolved in the clearance and endocytosis of hemolopion/haboratoollobic norvolved in the clearance and endocytosis of hemolopion/haboratoollobic norvolved in the clearance and endocytosis of hemolopion/haboratory experiessed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemolopion/haboratory experiessed.	Reactivity	Human,Mouse
Alternative Names M130; MM130; SCARI1 Research Field Immunology Product Categories Primary antibody Host Mouse Molecular Weight Calculated MW: 125 kDa Clonality Monoclonal Antibody Clonality No. 9H9-H6-W6 Dilution WB: 1/500-1/1000 IHC: 1/100-1/200 ELISA: 1/10000 Immunogen Purified recombinant fragment of human CD163 expressed in E. Coli. Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemodoloin/haptoclobin complexes by	Applications	WB,IHC-P,ELISA
Product Categories Primary antibody Host Mouse Molecular Weight Calculated MW: 125 kDa Clonality Monoclonal Antibody Clonality No. 9H9-H6-W6 Dilution WB: 1/500-1/1000 IHC: 1/100-1/200 ELISA: 1/10000 Immunogen Purified recombinant fragment of human CD163 expressed in E. Coli. Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The Description of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemodoloin/haptoclobin complexes by	Gene Name	CD163
Product Categories Primary antibody Host Mouse Molecular Weight Calculated MW: 125 kDa Clonality Monoclonal Antibody Clonality No. 9H9-H6-W6 Dilution WB: 1/500-1/1000 IHC: 1/100-1/200 ELISA: 1/10000 Immunogen Purified recombinant fragment of human CD163 expressed in E. Coli. Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemodolobin/haptoglobin complexes by	Alternative Names	M130; MM130; SCARI1
Host Mouse Molecular Weight Calculated MW: 125 kDa Clonality Monoclonal Antibody Clonality No. 9H9-H6-W6 Dilution WB: 1/500-1/1000 IHC: 1/100-1/200 ELISA: 1/10000 Immunogen Purified recombinant fragment of human CD163 expressed in E. Coli. Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemodolobin/habtoolobin complexes by	Research Field	Immunology
Molecular Weight Calculated MW: 125 kDa Clonality Monoclonal Antibody Clonality No. 9H9-H6-W6 Dilution WB: 1/500-1/1000 IHC: 1/100-1/200 ELISA: 1/10000 Immunogen Purified recombinant fragment of human CD163 expressed in E. Coli. Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/handoplobin complexes by	Product Categories	Primary antibody
Clonality Monoclonal Antibody Clonality No. 9H9-H6-W6 Dilution WB: 1/500-1/1000 IHC: 1/100-1/200 ELISA: 1/10000 Immunogen Purified recombinant fragment of human CD163 expressed in E. Coli. Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by	Host	Mouse
Clonality No. 9H9-H6-W6 Dilution WB: 1/500-1/1000 IHC: 1/100-1/200 ELISA: 1/10000 Immunogen Purified recombinant fragment of human CD163 expressed in E. Coli. Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by	Molecular Weight	Calculated MW: 125 kDa
Dilution WB: 1/500-1/1000 IHC: 1/100-1/200 ELISA: 1/10000 Immunogen Purified recombinant fragment of human CD163 expressed in E. Coli. Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by	Clonality	Monoclonal Antibody
Immunogen Purified recombinant fragment of human CD163 expressed in E. Coli. Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by	Clonality No.	9H9-H6-W6
Purification Affinity Purified Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by	Dilution	WB: 1/500-1/1000 IHC: 1/100-1/200 ELISA: 1/10000
Conjugation Unconjugated Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by	Immunogen	Purified recombinant fragment of human CD163 expressed in E. Coli.
Modification Unmodified Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by	Purification	Affinity Purified
Form Liquid Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by	Conjugation	Unconjugated
Buffer System Purified antibody in PBS with 0.05% sodium azide Concentration 1 mg/ml Purity ≥90% Storage Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by	Modification	Unmodified
Concentration 1 mg/ml Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by	Form	Liquid
Purity ≥90% Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by	Buffer System	Purified antibody in PBS with 0.05% sodium azide
Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by	Concentration	1 mg/ml
cycles. Background The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by	Purity	≥90%
cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by	Storage	·
	Background	cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658








	oxidative damage. This protein may also function as an innate immune sensor for bacteria and inducer of local inflammation. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd