



TRMT10C rabbit pAb

货号	BYab-17794
同位型	lgG
种属	Human;Mouse;Rat
应用	WB
靶点	TRMT10C
基因名称	TRMT10C MRPP1 RG9MTD1
蛋白名称	Mitochondrial ribonuclease P protein 1 (Mitochondrial RNase P protein 1) (EC 2.1.1) (HBV pre-S2 trans-regulated protein 2) (RNA (guanine-9-)-methyltransferase domain-containing protein 1) (Renal car
免疫原	Synthesized peptide derived from human TRMT10C
特异性	This antibody detects endogenous levels of TRMT10C at Human, Mouse,Rat
组成	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
来源	Polyclonal, Rabbit,IgG
稀释	WB 1:500-2000
纯化工艺	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
分子量	44kD
功能	Mitochondrial tRNA N(1)-methyltransferase involved in mitochondrial tRNA maturation . Component of mitochondrial ribonuclease P, a complex composed of TRMT10C/MRPP1, HSD17B10/MRPP2 and PRORP/MRPP3, which cleaves tRNA molecules in their 5'-ends . Together with HSD17B10/MRPP2, forms a subcomplex of the mitochondrial ribonuclease P, named MRPP1-MRPP2 subcomplex, which displays functions that are independent of the ribonuclease P activity . The MRPP1-MRPP2 subcomplex catalyzes the formation of N(1)-methylguanine and N(1)-methyladenine at position 9 (m1G9 and m1A9, respectively) in tRNAs; TRMT10C/MRPP1 acting as the catalytic N(1)-methyltransferase subunit . The MRPP1-MRPP2 subcomplex also acts as a tRNA maturation platform: following 5'-end cleavage by the mitochondrial ribonuclease P complex, the MRPP1-MRPP2 subcomplex enhances the efficiency of 3'-processing catalyzed by ELAC2, retains th
细胞定位	Mitochondrion matrix, mitochondrion nucleoid .
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)

Nanjing BYabscience technology Co.,Ltd





有关注意事项

Avoid repeated freezing and thawing!

使用建议

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd