



Phospho-Nrf2 (Ser40) Rabbit mAb

Catalog No	BYab-17816
Isotype	IgG
Reactivity	Human
Applications	WB,IHC-P,ICC/IF,IP,FC
Gene Name	NFE2L2
Alternative Names	NFE2L2; NRF2; Nuclear factor erythroid 2-related factor 2; NF-E2-related factor 2; NFE2-related factor 2; HEBP1; Nuclear factor; erythroid derived 2; like 2
Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody
Host	Rabbit
Molecular Weight	Calculated MW: 68 kDa; Observed MW: 90 kDa
Clonality	Monoclonal Antibody
Clonality No.	R03-4A1
Dilution	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/50 FC: 1/50-1/100
Immunogen	A synthesized peptide derived from human Phospho-Nrf2 (S40)
Purification	Affinity Chromatography
Conjugation	Unconjugated
Modification	Phosphorylated
Form	Liquid
Buffer System	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Concentration	1 mg/ml
Purity	≥90%
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Background	Transcription activator that binds to antioxidant response (ARE) elements in the promoter regions of target genes. Important for the coordinated up-regulation of genes in response to oxidative stress. May be involved in the transcriptional activation of genes of the beta-globin cluster by mediating enhancer activity of

Nanjing BYabscience technology Co.,Ltd



hypersensitive site 2 of the beta-globin locus control region.


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd