



ACOX1 rabbit pAb

货号	BYab-17982
同位型	IgG
应用	WB
种属	Human;Mouse;Rat
靶点	ACOX1
基因名称	ACOX1 ACOX
蛋白名称	Peroxisomal acyl-coenzyme A oxidase 1 (AOX) (EC 1.3.3.6) (Palmitoyl-CoA oxidase) (Straight-chain acyl-CoA oxidase) (SCOX)
免疫原	Synthesized peptide derived from human ACOX1
特异性	This antibody detects endogenous levels of ACOX1 at Human, Mouse,Rat
组成	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
来源	Polyclonal, Rabbit,IgG
稀释	WB 1:500-2000
纯化工艺	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
分子量	73kD
功能	Involved in the initial and rate-limiting step of peroxisomal beta-oxidation of straight-chain saturated and unsaturated very-long-chain fatty acids . Catalyzes the desaturation of fatty acyl-CoAs such as palmitoyl-CoA (hexadecanoyl-CoA) to 2-trans-enoyl-CoAs ((2E)-enoyl-CoAs) such as (2E)-hexadecenoyl-CoA, and donates electrons directly to molecular oxygen (O(2)), thereby producing hydrogen peroxide (H(2)O(2)) . ; [Isoform 1]: Shows highest activity against medium-chain fatty acyl-CoAs. Shows optimum activity with a chain length of 10 carbons (decanoyl-CoA) in vitro. ; [Isoform 2]: Is active against a much broader range of substrates and shows activity towards long-chain fatty acyl-CoAs.
细胞定位	Peroxisome .
组织表达	Widely expressed with highest levels of isoform 1 and isoform 2 detected in testis. Isoform 1 is expressed at higher levels than isoform 2 in liver and kidney while isoform 2 levels are higher in brain, lung, muscle, white adipose tissue and testis. Levels are almost equal in heart.
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)

Nanjing BYabscience technology Co.,Ltd



有关注意事项

Avoid repeated freezing and thawing!

使用建议

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images