



PARVB rabbit pAb

| | |
|--------|--|
| 货号 | BYab-18060 |
| 同位型 | IgG |
| 应用 | WB |
| 种属 | Human;Mouse |
| 靶点 | PARVB |
| 基因名称 | PARVB CGI-56 |
| 蛋白名称 | Beta-parvin (Affixin) |
| 免疫原 | Synthesized peptide derived from human PARVB |
| 特异性 | This antibody detects endogenous levels of PARVB at Human, Mouse |
| 组成 | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| 来源 | Polyclonal, Rabbit,IgG |
| 稀释 | WB 1:500-2000 |
| 纯化工艺 | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| 分子量 | 40kD |
| 功能 | Adapter protein that plays a role in integrin signaling via ILK and in activation of the GTPases CDC42 and RAC1 by guanine exchange factors, such as ARHGEF6. Is involved in the reorganization of the actin cytoskeleton and formation of lamellipodia. Plays a role in cell adhesion, cell spreading, establishment or maintenance of cell polarity, and cell migration. |
| 细胞定位 | Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cytoskeleton. Cell projection, lamellipodium. Cytoplasm, myofibril, sarcomere. Cytoplasm, myofibril, sarcomere, Z line. Constituent of focal adhesions. Detected at the tips of the leading edge of cells. Colocalizes with F-actin at the tips of lamellipodia. |
| 组织表达 | Expressed predominantly in heart and skeletal muscle. |
| 浓度 | 1 mg/ml |
| 储存 | -15°C to -25°C/1 year(Do not lower than -25°C) |
| 有关注意事项 | Avoid repeated freezing and thawing! |

Nanjing BYabscience technology Co.,Ltd



使用建议

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images